



AD 690185

AD

TECHNICAL REPORT

WVT-6923

MAXIMUM LIKELIHOOD ESTIMATION OF DISTRIBUTION PARAMETERS
FROM GENERALLY CENSORED SAMPLES

BY

ROYCE W. SOANES, JR

JUNE 1969

BENET R&E LABORATORIES

WATERVLIET ARSENAL

WATERVLIET—NEW YORK

REPRODUCED BY
CLEARINGHOUSE
FOR THE ARMY
AND NAVY
AND AIR FORCE

THIS REPORT IS THE PROPERTY OF THE ARMY
AND IS LOANED TO YOUR ORGANIZATION; IT
AND ITS CONTENTS ARE NOT TO BE DISTRIBUTED

/s

DISPOSITION

Destroy this report when it is no longer needed. Do not return it to the originator.

DISCLAIMER

The findings in this report are not to be construed as an official Department of the Army position.

MAXIMUM LIKELIHOOD ESTIMATION OF DISTRIBUTION PARAMETERS
FROM GENERALLY CENSORED SAMPLES

ABSTRACT

Cross-Reference Data

A simple derivation of Maximum Likelihood
the likelihood function for Likelihood Function
generally censored samples Censored Data
is given, where "generally Truncated Tests
censored sample" connotes:
a set of failure data con-
taining both failures and
survivors with various dif-
ferent tentative survival
times.

TABLE OF CONTENTS

<u>SECTION</u>		<u>PAGE</u>
	Abstract	1
1	Introduction	3
2	Multinomial Process	4
3	Likelihood Function	6
	References	14
	DD Forms 1473	

1. INTRODUCTION

In an ideal situation, one is presented with a set of failure data in which every piece of data indicates that a failure has taken place, i.e., one has information on failures and on failures only. In this simplest of situations, parameter estimation may be carried out in a variety of relatively simple ways.

More often than not, however, this situation is not the case; instead, one is presented with a number of indicated failures and some additional information on items which have been tested to varying degrees, but not to failure. The reason for lack of further testing on the unfailed specimens may be an economic one or it may be that the unfailed items have been discarded for virtually any reason. Irrespective of the reasons involved, the data has been "censored", i.e., someone has either purposely or inadvertently prevented us from seeing just exactly when some of the items tested will fail. This report deals with the general method of maximum likelihood as applied to this general censoring problem.

2. MULTINOMIAL PROCESS

The multinomial process is of central importance in the derivation of the censored likelihood function. A multinomial process is simply the repeated performance of an experiment E n times where the experiment has r (mutually exclusive and exhaustive) outcomes with associated probabilities. A description of the process may be abbreviated symbolically:

Perform experiment E n times where the outcomes of E are:

$$O_1, O_2, \dots, O_r$$

and

$$P(O_i) = p_i \quad i = 1, 2, \dots, r$$

Repetition of experiment E may be viewed as a composite experiment E_c , a typical outcome of which is:

$$O_c \equiv n_1 O_1', n_2 O_2', \dots, n_r O_r'$$

are observed. The probability of this typical outcome of E_c is given by:

$$P(Q_c) = P(n_1, n_2, \dots, n_r)$$

$$= \frac{n!}{n_1! n_2! \dots n_r!} p_1^{n_1} p_2^{n_2} \dots p_r^{n_r}$$

(1)

This formula is given in any elementary statistics text under the heading "multinomial distribution." It should be noted that $P(Q_c)$ is merely the probability of an outcome of E_c ; in general one wants the probability of an event (aggregation of outcomes). An event is given by a set of conditions on the n_i .

Therefore,


$$P(A) = \sum_{\substack{\text{(such that the } n_i \text{ satisfy} \\ \text{condition A)}}} P(n_1, n_2, \dots, n_r)$$

(2)

3. LIKELIHOOD FUNCTION

The principle of maximum likelihood dictates that we find the parameters which maximize the theoretical probability of obtaining the results we actually did obtain, i.e., other results "should" be less likely than the actual results.

The theoretical probability of obtaining the observed results must therefore be obtained first. The event describing the observed results is: K x 's fail at specific times x_1, x_2, \dots, x_K and N x 's survive for various lengths of time $s_1 \leq s_2 \leq \dots \leq s_N$ without failing ($N + K = n$). Pictorially, the actual test results will look like:



A theoretical one by one test of each item to failure may be viewed as a multinomial process with the following outcomes (for E).

$$Q'_1 \equiv x \in (x_1, x_1 + dx)$$

$$Q'_2 \equiv x \in (x_2, x_2 + dx)$$

⋮

$$Q'_K \equiv x \in (x_K, x_K + dx)$$

$$Q_1 \equiv x \in (x_1, x_2)$$

$$Q_2 \equiv x \in (x_2, x_3)$$

⋮

$$Q_{N-1} \equiv x \in (x_{N-1}, x_N)$$

$$Q_N \equiv x \in (x_N, \infty)$$

$$Q_{N+1} \equiv x \text{ is elsewhere.}$$

Strictly speaking, these outcomes are not mutually exclusive in a set theoretic sense; they are in a measure theoretic sense, however, and this is sufficient. This abuse of notation should be tolerated here for the sake of brevity.

If f is the underlying density function and F is the distribution function, the probabilities corresponding to these outcomes are:

$$P'_1 = f(x_1) dx$$

$$P'_2 = f(x_2) dx$$

$$\vdots$$

$$P'_K = f(x_K) dx$$

$$P_1 = F(s_2) - F(s_1)$$

$$P_2 = F(s_3) - F(s_2)$$

$$\vdots$$

$$P_{N-1} = F(s_N) - F(s_{N-1})$$

$$P_N = F(\infty) - F(s_N) = 1 - F(s_N)$$

Now that E has been specified, the probability of obtaining the observed results in a theoretical test of N items to failure is desired. The event describing the observed results is:

$$n'_1 = 1$$

$$n'_2 = 1$$

$$\vdots$$

$$n'_K = 1$$

$$n_1 = k_1$$

$$n_2 = k_2$$

$$\vdots$$

$$n_N = k_N$$

$$n_{N+1} = 0$$

and

$$k_i \leq 1$$

$$k_1 + k_2 \leq 2$$

$$\vdots$$

$$\sum_{i=1}^j k_i \leq j$$

$$\vdots$$

$$\sum_{i=1}^N k_i = N$$

The probability of obtaining the observed results is therefore:

$$P = \sum \frac{n!}{1!1!\dots1!k_1! \dots k_N!} p_1' \dots p_K' p_1^{k_1} \dots p_N^{k_N} p_{N+1}^0$$

$$(l \leq N \Rightarrow \sum_{i=1}^l k_i \leq l,)$$

$$\sum_{i=1}^N k_i = N$$
(3)

$$= \sum \frac{n!}{k_1! \dots k_N!} \left[\prod_{i=1}^K f(x_i) dx \right] \left[\prod_{j=1}^N \{F(x_{j+1}) - F(x_j)\}^{k_j} \right]$$

$$(l \leq N \Rightarrow \sum_{i=1}^l k_i \leq l,)$$

$$\sum_{i=1}^N k_i = N$$
(4)

where $x_{N+1} = \infty$

These sums are to be taken over all k such that

$$\sum_{i=1}^l k_i \leq l \quad (\text{for all } l \leq N)$$

That is, for each term of the sum, all k 's must be assigned and they must be assigned such that for all $l \leq N$,

$$\sum_{i=1}^l k_i \leq l.$$

The symbol



denotes "implies."

The likelihood function is therefore defined:

$$L = \sum \frac{n!}{k_1! \dots k_N!} \left[\prod_{i=1}^K f(x_i) \right] \left[\prod_{j=1}^N \left\{ F(a_{j+1}) - F(a_j) \right\}^{k_j} \right]$$

$$(l \leq N \Rightarrow \sum_{i=1}^l k_i \leq l, \quad (a_{N+1} = \infty))$$

$$\sum_{i=1}^N k_i = N$$

(5)

This expression may be written logarithmically:

$$L = \sum \frac{n!}{k_1! \dots k_N!} \exp \left\{ \sum_{i=1}^K \ln f(x_i) + \sum_{j=1}^N k_j \ln [F(a_{j+1}) - F(a_j)] \right\}$$

$$(l \leq N \Rightarrow \sum_{i=1}^l k_i \leq l, \quad \sum_{i=1}^N k_i = N)$$

(6)

Equation (5) may be written more succinctly if one of the products is factored out:

$$L = \prod_{i=1}^K f(x_i) \sum_{\substack{k_1, \dots, k_N \\ \left(\begin{array}{l} l \leq N \Rightarrow \sum_{i=1}^l k_i \leq l, \\ \sum_{i=1}^N k_i = N \end{array} \right)}} \frac{n!}{k_1! \dots k_N!} \prod_{j=1}^N \left\{ F(x_{j+1}) - F(x_j) \right\}^{k_j}$$

(7)

$$= S \prod_{i=1}^K f(x_i)$$

Consider the case when say $N = 3$. There are five terms for S in this case; the k 's for each term are given by the table:

	k_1	k_2	k_3
S_1	1	1	1
S_2	0	2	1
S_3	1	0	2
S_4	0	1	2
S_5	0	0	3

An important special case arises when $a_1 = a_2 =$
 $= a_3 = \dots = a_N$. In this case, all terms of
 S vanish except one; the maximization of the likeli-
hood function is therefore considerably easier. The like-
lihood function will in general be amenable to maximization
only through an approximate numerical method. Such a
method may involve taking the first or second partial
derivatives of L with respect to the parameters.

REFERENCES:

Shooman, M. L., Probabilistic Reliability,
McGraw-Hill, pp. 475 - 480

DOCUMENT CONTROL DATA - R & D

(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)

1. ORIGINATING ACTIVITY (Corporate author) Watervliet Arsenal Watervliet, N.Y. 12189		2a. REPORT SECURITY CLASSIFICATION Unclassified	
		2b. GROUP	
3. REPORT TITLE MAXIMUM LIKELIHOOD ESTIMATION OF DISTRIBUTION PARAMETERS FROM GENERALLY CENSORED SAMPLES			
4. DESCRIPTIVE NOTES (Type of report and inclusive dates) Technical Report			
5. AUTHOR(S) (First name, middle initial, last name) Royce W. Soanes, Jr.			
6. REPORT DATE June 1969		7a. TOTAL NO. OF PAGES 16	7b. NO. OF REFS 1
8a. CONTRACT OR GRANT NO. -		9a. ORIGINATOR'S REPORT NUMBER(S) WVT-6923	
b. PROJECT NO. -			
c.		9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)	
d.			
10. DISTRIBUTION STATEMENT This document has been approved for public release and sale; its distribution is unlimited.			
11. SUPPLEMENTARY NOTES		12. SPONSORING MILITARY ACTIVITY U. S. Army Weapons Command	
13. ABSTRACT A simple derivation of the likelihood function for generally censored samples is given, where "generally censored sample" connotes: a set of failure data containing both failures and survivors with various different tentative survival times.			

14.	KEY WORDS	LINK A		LINK B		LINK C	
		ROLE	WT	ROLE	WT	ROLE	WT
	Maximum Likelihood						
	Likelihood Function						
	Censored Data						
	Truncated Tests						

AD Accession No.
Benet Laboratories, Watervliet Arsenal, Watervliet, N.Y.
MAXIMUM LIKELIHOOD ESTIMATION OF DISTRIBUTION PARAMETERS
FROM GENERALLY CENSORED SAMPLES by Royce W. Soanes, Jr.
Report No. WWT-6923, June 1969, 16 pages. Unclassified
Report

A simple derivation of the likelihood function for generally censored samples is given, where "generally censored sample" connotes: a set of failure data containing both failures and survivors with various different tentative survival times.

Maximum Likelihood
Likelihood Function
Censored Data
Truncated Tests

Distribution
Unlimited

AD Accession No.
Benet Laboratories, Watervliet Arsenal, Watervliet, N.Y.
MAXIMUM LIKELIHOOD ESTIMATION OF DISTRIBUTION PARAMETERS
FROM GENERALLY CENSORED SAMPLES by Royce W. Soanes, Jr.
Report No. WWT-6923, June 1969, 16 pages. Unclassified
Report

A simple derivation of the likelihood function for generally censored samples is given, where "generally censored sample" connotes: a set of failure data containing both failures and survivors with various different tentative survival times.

Maximum Likelihood
Likelihood Function
Censored Data
Truncated Tests

Distribution
Unlimited

AD Accession No.
Benet Laboratories, Watervliet Arsenal, Watervliet, N.Y.
MAXIMUM LIKELIHOOD ESTIMATION OF DISTRIBUTION PARAMETERS
FROM GENERALLY CENSORED SAMPLES by Royce W. Soanes, Jr.
Report No. WWT-6923, June 1969, 16 pages. Unclassified
Report

A simple derivation of the likelihood function for generally censored samples is given, where "generally censored sample" connotes: a set of failure data containing both failures and survivors with various different tentative survival times.

Maximum Likelihood
Likelihood Function
Censored Data
Truncated Tests

Distribution
Unlimited

AD Accession No.
Benet Laboratories, Watervliet Arsenal, Watervliet, N.Y.
MAXIMUM LIKELIHOOD ESTIMATION OF DISTRIBUTION PARAMETERS
FROM GENERALLY CENSORED SAMPLES by Royce W. Soanes, Jr.
Report No. WWT-6923, June 1969, 16 pages. Unclassified
Report

A simple derivation of the likelihood function for generally censored samples is given, where "generally censored sample" connotes: a set of failure data containing both failures and survivors with various different tentative survival times.

Maximum Likelihood
Likelihood Function
Censored Data
Truncated Tests

Distribution
Unlimited

AD <u>Benet Laboratories, Watervliet Arsenal, Watervliet, N.Y.</u> MAXIMUM LIKELIHOOD ESTIMATION OF DISTRIBUTION PARAMETERS FROM GENERALLY CENSORED SAMPLES by Royce W. Soanes, Jr. Report No. WWT-6923, June 1969, 16 pages. Unclassified Report	AD <u>Benet Laboratories, Watervliet Arsenal, Watervliet, N.Y.</u> MAXIMUM LIKELIHOOD ESTIMATION OF DISTRIBUTION PARAMETERS FROM GENERALLY CENSORED SAMPLES by Royce W. Soanes, Jr. Report No. WWT-6923, June 1969, 16 pages. Unclassified Report	Maximum Likelihood Likelihood Function Censored Data Truncated Tests Distribution Unlimited	Maximum Likelihood Likelihood Function Censored Data Truncated Tests Distribution Unlimited
AD <u>Benet Laboratories, Watervliet Arsenal, Watervliet, N.Y.</u> MAXIMUM LIKELIHOOD ESTIMATION OF DISTRIBUTION PARAMETERS FROM GENERALLY CENSORED SAMPLES by Royce W. Soanes, Jr. Report No. WWT-6923, June 1969, 16 pages. Unclassified Report	AD <u>Benet Laboratories, Watervliet Arsenal, Watervliet, N.Y.</u> MAXIMUM LIKELIHOOD ESTIMATION OF DISTRIBUTION PARAMETERS FROM GENERALLY CENSORED SAMPLES by Royce W. Soanes, Jr. Report No. WWT-6923, June 1969, 16 pages. Unclassified Report	Maximum Likelihood Likelihood Function Censored Data Truncated Tests Distribution Unlimited	Maximum Likelihood Likelihood Function Censored Data Truncated Tests Distribution Unlimited